

MONTANA TOBACCO USE PREVENTION PROGRAM

## PROGRESS REPORT

JULY 2010 - JUNE 2012

## **Mission & Goals**

The mission of the Montana Tobacco Use Prevention Program (MTUPP) is to address the public health crisis caused by the use of all forms of commercial tobacco products. MTUPP is working to eliminate tobacco use, particularly among young people, via programs and policies throughout Montana.

- Preventing initiation among youth
   and young adults Most people who start
   smoking are younger than 18 years of age, making
   intervening critical. Community programs and
   school-based policies and interventions should
   be part of a comprehensive effort, implemented
   in coordination across the community and school
   environments and in conjunction with increasing
   the unit price of tobacco products, sustaining anti tobacco media campaigns, making environments
   smoke-free, and engaging in other efforts to
   create tobacco-free social norms.
- Promoting quitting among adults and youth — Interventions that increase quitting can decrease premature mortality and tobacco-related healthcare costs in the short-term.
- Eliminating exposure to secondhand smoke — Secondhand smoke causes premature death and disease in children and adults who do not smoke. There is no risk-free level of exposure to secondhand smoke.
- Identifying and eliminating tobaccorelated disparities among population groups — Some populations experience a disproportionate health and economic burden from tobacco use; therefore, a focus on eliminating such tobacco-related disparities is necessary.

## From the Director



Anna Whiting Sorrell

Please stop and consider how tobacco addiction has affected your life and the lives of your family and friends. Few Montanans can say they have never known someone whose life has not been altered by commercial tobacco—be it a loss

of finances, death from lung cancer or heart disease, or the premature passing of a loved one. The diseases caused by tobacco addiction are still the leading causes of death in America and in Montana, claiming the lives of an average of four Montanans each day.

Tobacco addiction complicates many other serious chronic diseases such as diabetes, hypertension, asthma and mental illness. MTUPP is leading the cooperative effort to ensure that tobacco prevention and cessation are addressed throughout the state, providing a comprehensive money-saving approach and health improvement.

Please join me in thanking our partners, including our team of dedicated tobacco prevention specialists across Montana, by recognizing their accomplishments and supporting the important work being done in our communities. As you will see, this is a multi-faceted program and we invite you to join our efforts.

ausoneel


**Anna Whiting Sorrell**, *Director*,
Department of Public Health and Human Services




Tobacco use is the leading cause of preventable death and disease in Montana. Tobacco use causes many deadly diseases, including cardiovascular disease and cancer, the leading causes of death in Montana and the United States. An estimated 1,400 Montanans die from diseases caused by tobacco use every year.1



## Number of Tobacco Related Deaths by County, 2004-2009, Montana




## Dollars Spent by Tobacco Companies on Marketing & Advertising vs. Dollars Spent by the State of Montana on Tobacco Prevention<sup>2</sup>




To counter the impact that Clean Indoor Air Acts have on cigarette sales, tobacco companies have increased their line of smokeless tobacco products—particularly products appealing to youth. Marketing for smokeless tobacco in the US rose 400% from 1986 to 2006. It rose another 54% to almost \$548 million from 2006 to 2008. Tobacco has become even more addictive due to increased nicotine content of tobacco products.<sup>4</sup>

## Per Capita Cigarette Packs Sold in Montana, 1955-2011<sup>3</sup>



# SPOTE GETT Quitting Tobacco


- Montanans are using this free service and successfully quitting.
   There have been over 62,000 calls since 2004.
- The Montana Tobacco Quit Line ranks in the top 10 in the country for comprehensive services provided.<sup>6</sup>
- The success rate for quitting is 40% higher for participants who receive both coaching and Chantix than for those who try to quit on their own. <sup>7</sup>



## SPOTE STATE OF THE STATE OF THE


- Nearly 90% of tobacco users started before the age of 18, and 99% started before the age of 26.8 If all people under the age of 26 could be persuaded never to try any tobacco product, nearly 18,000 Montana youth could be spared an eventual death caused by tobacco-related illness.9
- All 60 MTUPP funded community and youth programs have a youth coalition
  affiliated with reACT Against Corporate Tobacco, who work to change the
  social norm so that our Montana youth never use tobacco.
- reACT also offers mini-grants across the state to local youth coalitions.
   Grant activities reached an estimated 5,000 teens and 10,000 adults across the state in FY 2012.

## Prevalence of Tobacco Use Among Montana Youth from 1995-2011, YRBS



**ABOVE** Smoking prevalence among Montana youth has been steadily declining since 1997. Smokeless tobacco use among this population has not significantly changed in the past 10 years.

## Schools District with Comprehensive Tobacco Free School Policy as of June 2012



**ABOVE** MTUPP partners with the Office of Public Instruction to increase the number of Montana school districts who adopt Comprehensive Tobacco Free School Policy and provides media literacy training. As of June 2012, a total of 194 school districts (46%) have adopted Comprehensive Tobacco Free School Policy, compared to 23 school districts as of September 2010.

RIGHT Montana's youth tobacco prevention program, reACT, has partnered with the Montana High School Rodeo Association (MHSRA). As a result, in June 2011, MHSRA passed a comprehensive tobacco free policy, which prohibits tobacco use of any kind at all MHSRA sanctioned events.



## COMMUNITY Partners Involved CITIZINS



of Montanans are served by tobacco use prevention programs located in 50 counties, on seven American Indian reservations and two Urban Indian Centers.

### Youth

- Youth smoking rates declined from 29% in 2001 to 17% in 2011. 10
- 30% of Montana high school senior boys are current smokeless tobacco users.

### Secondhand Smoke Exposure

- Secondhand smoke exposure, even for as little as 30 minutes, can cause heart attacks. People at risk for heart disease or stroke are particularly vulnerable. 12
- 11% of adults were exposed to secondhand smoke at home<sup>14</sup> and 33% of children were exposed while they were riding in cars. <sup>14</sup>



## **Quitting Tobacco**

- People who use the quit line are 7–10 times more likely to successfully quit.
- 63% of tobacco users want to quit. 16

## **Tobacco-related disparities**

- In 2010, the smoking rate for American Indian adults was 49%<sup>17</sup>
- 44% of Montanans in the lowest income bracket (<\$15,000) are current smokers compared to only 7% of Montanans in the highest income bracket (\$75,000+).

## SPOTELSHIT Specific Populations

- 17% of Montana mothers smoked during their pregnancy 19
- Prenatal exposure to tobacco smoke is associated with premature births, putting babies at risk for serious health problems and drastically increasing medical costs.

### **Montana Medicaid benefits**

Tobacco dependence treatment is now more accessible than ever for clients of Montana Medicaid. They can access counseling from a wide variety of health professionals and licensed counselors and receive all FDA approved cessation medications. MTUPP and Montana Medicaid are working together to ensure that all clients are aware of these services. Medicaid clients comprised more than 10% of Montana Quit Line callers this past year.<sup>21</sup>





## Consumers with Mental Illness or Substance Abuse Disorders

People with mental illness or addictions have some of the highest rates of tobacco use in the United States; people with psychiatric disorders smoke at rates almost twice as high as the general population (41% versus 22.5%, respectively). <sup>23</sup> MTUPP partnered with behavioral health service providers and consumers to reach this often-neglected group.

Site visits to outpatient treatment facilities helped raise staff and client awareness of the toll of tobacco on this disparate population and how tobacco use can interfere with recovery. This outreach included creating specialized health promotion materials, providing expert training for health professionals, fostering peer support programs, and offering assistance with the development of tobacco free policies and treatment options.



## **American Indians**

During February 2010 to June 2012, MTUPP met its goal of doubling American Indian callers to the guit line. Montana's seven tribes and two Urban Indian Centers were featured in an aggressive media campaign aimed at increasing knowledge of secondhand smoke and awareness of the quit line.

> Get FREE help quitting tobacco today. he Montana Tobacco Quit Line offers:

A FREE personalized quit plan

· FREE educational materials FREE phone coaches

Quit Tobacco Today!

1-800-QUIT-NOW



cancer, heart disease and other major health problems. But women who smoke during pregnancy put themselves and their unborn babies at risk for other health problems. The dangers of smoking during pregnancy include premature birth, certain birth defects and infant death 22

Babies exposed to secondhand smoke are more likely to die from SIDS (Sudden Infant Death Syndrome). 20 Average first-year medical costs are about 10 times greater for preterm infants than for full-term infants. 20

## SPOTE SECOND Second hand Smoke

Exposure to secondhand smoke increases the risk of health problems:

- Heart disease, increased heart rate, increased blood pressure and heart attacks
- Lung cancer, emphysema, bronchitis, asthma attacks and pneumonia
- Ear infections, throat, nose and eye irritation

Secondhand smoke knows no boundaries. Smokefree policies ensure healthier living for everyone:

- Protect all people
- Reduce youth initiation and change social norms
- Encourage quitting

Passing policies that prohibit smoking (smokefree policies) and the use of smokeless tobacco products as well (tobacco free policies) actively change the social norms related to tobacco use. They also create an environment that incentivizes and encourages tobacco users to quit. Studies have shown that challenging the perception of smoking as a normal adult behavior through smokefree policies can change the attitudes and behaviors of adolescents. This can result in a reduction of the number of adolescents who start using tobacco.

Montana continues to work closely with medical campuses, college campuses and multi-unit housing owners and managers to encourage the adoption of smokefree and tobacco free policies. The map on the following page shows progress made over the last seven years. During the last two years alone, 32 new policies have been adopted across Montana, including those on 18 medical campuses, seven college campuses and seven housing authorities.

## Tobacco Free & Smokefree


## POLICY MAPS

*2005* 



- Medical Campuses (Tobacco Free/Smokefree)
- Housing Authorities (Smokefree)
- College Campuses (Tobacco Free)

2012



### **Medical Campuses**

#### **Tobacco Free Policies**

Barrett Hospital and Health Care, Dillon BCBS of Montana, Helena

Big Sandy Medical Center, Big Sandy

Broadwater Health Center, Townsend

Butte Community Health Clinic, Butte Central Montana Medical Center. Lewiston

Clark Fork Valley Hospital, Plains

Community Health Partners, Livingston

Community Hospital of Anaconda, Anaconda

Community Medical Center, Missoula

Daniel's Memorial Health Care Center, Scobey

Discovery Care Center, Hamilton

Gallatin Rest Home, Bozeman

Glen-wood Inc., Plentywood

Glendive Medical Center, Glendive

Holy Rosary Healthcare, Miles City

Kalispell Regional Medical Center, Kalispell

Lewis and Clark County Health Department, Helena

Liberty County Hospital, Chester

Livingston Health Care, Livingston

Madison Valley Hospital-Clinic, Ennis

Marcus Daly Memorial Hospital, Hamilton

Marias Medical Center, Shelby

McCone County Hospital, Circle

Mineral Community Hospital, Superior

Missoula Health Department, Missoula

Missouri River Medical Center, Fort Benton

Montana Chemical Dependency Center, Butte

Montana Development Center, Boulder

Montana Mental Health Nursing Care Center, Lewistown

Montana State Hospital, Warm Springs

North Valley Hospital, Whitefish

Northern Montana Healthcare, Havre

Phillips County Hospital, Malta

Powell County Medical Center, Big Timber

Rosebud Health Care Center, Forsyth

Saint Joseph Hospital, Polson

Saint Luke Community Hospital, Ronan

Shephard's Way Assisted Living, Lewistown

Sheridan Memorial Hospital, Plentywood

Shodair Children's Hospital, Helena

Sidney Health Center, Sidney

St. James Hospital, Butte

St. John's Lutheran Hospital, Libby

St. Patrick Hospital, Missoula

St. Pete's Medical Center, Helena

St. Peter's Hospital, Helena

Stillwater Community Hospital, Columbus

Tamarack Management, Missoula

Teton County Rest Home, Choteau

Wheatland Memorial Healthcare, Harlowton

#### **Smokefree Policies**

Benefis Health Care, Great Falls Billings Clinic, Billings Bozeman Deaconess Hospital, Bozeman St. Vincent's Health Care, Billings

### **Housing Authorities**

## Comprehensive Campus Wide or Facility Wide Smokefree Policies

Dawson County Housing Authority, Glendive Helena Housing Authority, Helena Housing Authority of Billings, Billings Miles City Housing Authority, Miles City Missoula Housing Authority, Missoula Richland County Housing Authority, Sidney Whitefish Housing Authority, Whitefish

#### **Partial Facility Smokefree Policies**

Great Falls Housing Authority, Great Falls Housing Authority of Butte, Butte Ronan Housing Authority, Ronan

### **College Campuses**

#### **Tobacco Free Policies**

Fort Peck Community College, Wolf Point Montana State University, Bozeman Montana State University, Billings Montana Tech, Butte University of Montana, Helena College of Technology, Helena University of Montana, Missoula University of Montana Western, Dillon





#### SOURCES

- CDC. Smoking-Attributable Mortality, Morbidity and Economic Costs (SAMMEC). Available at http://apps.nccd.cdc.gov/sammec/index.asp
- Campaign for Tobacco Free Kids state-specific tobacco company marketing expenditures 1998-2008. http://www.tobaccofreekids.org/ research/factshseets/pdf/0271.pdf. Accessed September 2010
- Orzechowski and Walker. The Tax Burden on Tobacco, Historical Compilation. Vol. 46, 2011
- 4) The Massachusetts Tobacco Control Program, Massachusetts Department of Public Health, Change in Nicotine Yields 1998 – 2004, Lois Keithly, Ph.D., Director; Doris Cullen, MA, Research Analyst, Thomas Land, Ph.D., Research Analyst.
- 5) Montana Tobacco Use Prevention Program Quit Line records: 2004-2012 (June)
- 6) American Lung Association, State of Tobacco Control, 2012
- 7) National Jewish Health Tobacco Cessation Outcome Results, Montana Tobacco Use Prevention Program, 2011

- 8) Preventing Tobacco Use Among Youth and Young Adults: A Report of the Surgeon General, 2012
- DPHHS State Improvement Plan Presentation, Billings, May 2012.
- 10) Montana Youth Risk Behavior Survey, 2011
- 11) ibid
- 12) The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General, 2006
- 13) Montana Department of Public Health and Human Services, Montana Tobacco Use Prevention Program. Montana Adult Tobacco Survey, 2004-2006, 2008-2009
- 14) Montana Department of Public Health and Human Services, Addictive and Mental Disorders Division. Prevention Needs Assessment, 2000-2008
- 15) American Cancer Society, Guide to Quitting Smoking, www. Cancer.org/Healthy/ StayAwayFromTobacco/Guide –to-Quitting-Smoking/guide-to-quitting-smoking-successrates. Accessed September 2012.

- CDC http://www.cdc.gov.tobacco/data\_ statistics/fact\_sheets/fast\_facts/. Accessed September 2012.
- 17) Montana Behavioral Risk Factor Surveillance System, 2010
- 18) Montana Birth Certificate data, Montana Office of Vital Statistics, 2008-2010
- Tobacco Control Legal Consortium, Cause and Effect: Tobacco Marketing Increases Youth Tobacco Use, Findings from the 2012 Surgeon General's Report (2012).
- 20) March of Dimes Foundation. http:// marchofdimes.com/pregnancy/alcohol\_ smoking.html. Accessed September 2011
- 21) Montana Tobacco Use Prevention Program
  Quit Line records
- 22) Centers for Disease Control and Prevention,
  Division of Reproductive Health
- 23) Karen Lasser et al., "Smoking and Mental Illness: A Population-Based Prevalence Study." The Journal of American Medical Association 284:20 (2000): 2606-2610.

#### **TOBACCO ADVISORY BOARD**

Robin Morris, Chair President, Morris & Morris, Inc., Columbus

Richard Sargent, MD, Vice-Chair Family Physician, St. Peter's Hospital, Helena

John Beck, LCPC
Director, Mental Health Center,
Livingston

Brett Christiansen Montana Collegiate Tobacco Prevention Initiative, Dillon

Rick Gale Tobacco Prevention Specialist, Gallatin County Ron Gleason CEO, Liberty Medical Center, Liberty County

Cristen Morigeau-Two Teeth Tobacco Prevention Specialist, Confederated Salish & Kootenai Tribes

Kristin Page-Nei Director, ACS State Government Relations, Missoula

Jan Parmelee
Tobacco Prevention Specialist,
Sanders County

Sharon Patton-Griffin, Associate Principal North Middle School, Great Falls Melanie Reynolds Health Officer, Lewis & Clark County

Marianne Roose Commissioner District 3, Lincoln County

Jeanne Seifert Health Officer, Dawson County

Robert Shepard, MD Family Physician, Helena

Marjorie Spotted Bird Tribal Health Director, Fort Peck



1,000 copies of this public document were published at an estimated cost of \$1.07 per copy, for a total cost of \$1,070.00, which includes \$1,070.00 for printing and \$157.00 for distribution. 1,000 copies of this public document were published at an estimated cost of \$1.07 per copy, for a total cost of \$1,070.00, which includes \$1,070.00 for printing and \$157.00 for distribution.